

- Progesterone in peripheral blood during the ovulatory cycle of the hen. *J. Endocrinol.* 57: 159
8. Koch, J.M., Moritz, J.S., Lay, D.C and Wilson, M.E(2005a). Motivation of hens to obtain feed during a molt induced by feed withdrawal, wheat middlings or melengesterol acetate. *Poult.Sci* 86:614-620
 9. Ladda, S., Bagliol, L., Lemi, G and Naitana, S(1999). Production and lambing rate of blatocyst derived from invitro matured oocyst after gonadotropin treatment of prepubertal ewes. *J. Anim. Sci.*, 77:2234-2239
 10. Parkhurst, C.R and Mountney, G.J (1988) Poultry meat and egg production. Page 203-205. C.R Parkhurst and G.J Mountney, eds. Chapman and Hall, New York.
 11. Richards, J. S. (1979). Hormonal control of ovarian follicular development: a 1978 perspective. *Rec. Prog. Hormone Res.*, 35:343.
 12. Steel, G.O and Torrie J.H (1980) Principles and procedure of statistics. McGraw Hill, New York.
 13. Sugano, M., Shinogi, T., Nakada, K and Nakao, T. (2001). Endocrine profile and embryo quality in Japanese black cattlesuperovulated with human menopausal gonadotropin and porcine follicle stimulating hormone. *Reproduction in Domestic Animals*. Volume 36(29)57